



Salamander Chytrid Disease

Other names: *Bsal*, Chytridiomycosis

- ♦ Salamander Chytridiomycosis, which is caused by the pathogen *Batrachochytrium salamandrivorans (Bsal)*, has spread from Asia to Europe where it has caused mass declines in salamander and newt populations.
- ♦ The international trade of amphibians and ornamental fish could facilitate the spread of the fungus to North America which is a biodiversity hotspot for salamanders and newts.

What is *Batrachochytrium salamandrivorans (Bsal)*?

- ♦ *Bsal* is a fungus that infects salamanders and newts (*Caudata*).
- ♦ The disease is similar to *B. dendrobatidis (Bd)*, a fungus which mostly infects frogs (*Anura*).
- ♦ *Bsal* has aquatic zoospores that infect the skin layers of an animal and can lead to skin lesions, anorexia, lethargy, ataxia and death.
- ♦ Common skin symptoms are similar to a bacterial infection: reddening, ulceration, and major skin shedding.
- ♦ The fungi can be spread through organic materials: mud, water, leaf litter, etc.
- ♦ Once introduced, the expectation is that *Bsal* is likely to survive and persist in most of Canada and will be impossible to eradicate.

What can you do to protect wild amphibians?

Observe and report

- ♦ Consider carrying a small jar of 70% ethanol in the field with your equipment and collecting fresh carcasses if you find dead salamanders or newts.
- ♦ Record the date and location of the observation and contact the **Canadian Wildlife Health Cooperative National Office [1(306) 966-5099]**.
- ♦ Do not freeze specimens. Preserving them in ethanol is best. If ethanol is not available, store specimens in a fridge for a short period of time.
- ♦ Take close-up photographs of the amphibians and send your pictures to the **CWHC (national@cwhc-rcsf.ca)**.


Take preventative measures

- ♦ Always follow a disinfection protocol to clean your field gear, whether you've worked at one or multiple sites (see page 2 for details).
- ♦ Do not move salamanders, newts or any other animals between sites.
- ♦ Captive salamanders, newts or other animals should never be released into the wild.
- ♦ Take appropriate steps to prevent disease spread with your captive salamanders. Follow good hygiene practices and quarantine abnormal-looking amphibians.

Have a look...



Fire salamander with lesions on its face. Photo by: Frank Pasmans, Gent University (2015).



Fire salamander with ulcers on its back. Photo by: Mark Blooi (2016)



Have a basic disinfection protocol ready!

Before leaving for the field, stock up on the following items:

- ◆ Brushes for scrubbing, antiseptic alcohol wipes, and bleach to disinfect.
- ◆ Handheld spray bottles and/or pump sprayers.
- ◆ Gloves for handling the animals. Use a new pair of gloves when handling individuals that would not otherwise come in contact with each other.
- ◆ Plastic containers or resealable bags of different sizes if you are collecting amphibians. Use a new container/bag for individuals that would not otherwise come into contact with each other.
- ◆ Bring additional sets of equipment if sampling in multiple locations (ideally one set per site).

After each sampling event and before moving on to the next site:

- ◆ Field gear (boots, clothes, boats, tires, etc.) should all be cleaned with soapy water and a brush to remove any excess organic material. Rinse the equipment with water (preferably hot tap water, if available).
- ◆ After washing and rinsing, apply a 5% bleach solution by spraying or immersing the gear: use $\frac{1}{2}$ cup of bleach for every $9\frac{1}{2}$ cups of water. If using or disposing of a large volume of bleach, do so away from all bodies of water or amphibian habitats. View the Canadian Wildlife Health Cooperative [Decontamination Protocol](#) for more details.
- ◆ If possible, rinse the bleached items with clean water (tap or bottle) to prevent exposing the next wetland to residual bleach and to minimize damage to the equipment. If in the field, rinse with water from the next sampling site. Allow the equipment to dry completely if you are done for the day.
- ◆ Use alcohol wipes to disinfect calipers, measuring boards, and other sensitive equipment.

1.



2.



3.



Images from: Northeastern Parc Partners in Amphibian and Reptile Conservation (2016) - Disinfection protocol.

For more information

- ◆ **Canadian Wildlife Health Cooperative:**
www.cwhc-rcsf.ca
- ◆ **The Amphibian Survival Alliance and Amphibian Specialist Group:**
www.amphibians.org
- ◆ **Documentary: The Origins of the Salamander Eater**
www.amphibians.org/news/watching-extinction-happen-origins-of-the-salamander-eater
- ◆ **The Chytrid disease website:**
www.salamanderfungus.org/about-bsal
- ◆ **The Chytrid disease Facebook Page:**
www.facebook.com/salamanderfungus



Top border photos courtesy of Crowley (2016): Northern Two-lined Salamander, Spotted Salamander, Red-spotted Newt, Blue-spotted Salamander, Northern Dusky Salamander, Eastern Red-backed Salamander, Four-toed Salamander, Northern Long-toed Salamander, Red-spotted Newt (aquatic). Bottom photo courtesy of Ovaska (2016).

 Environment and Climate Change Canada

Environnement et Changement climatique Canada

CANADIAN WILDLIFE HEALTH COOPERATIVE

CANADIAN DIRECTORS COMMITTEE COMITÉ DES DIRECTEURS CANADIENS DE LA FAUNE

 BRITISH COLUMBIA

Alberta Government

Manitoba

Ontario

Québec

New Brunswick

NOVA SCOTIA

Prince Edward Island

Communities, Land and Environment

Government of Northwest Territories

Gouvernement des Territoires du Nord-Ouest

Yukon

Nature CANADA

ASG

Canadian Herpetological Society

NWDC

THE NORTHEAST WILDLIFE DISEASE COOPERATIVE

Follow the CWHC:
   

Version: April 2017